

Polyamorphic Transformations in Fe-Ni-C Liquids: Implications for Chemical Evolution of Terrestrial Planets

Xiaojing Lai^{1,2*}, Bin Chen^{2*}, Jianwei Wang³, Yoshio Kono⁴, Feng Zhu⁵

¹Department of Geology and Geophysics, University of Hawaii at Manoa, Honolulu, Hawaii 96822, USA

²Hawaii Institute of Geophysics and Planetology, University of Hawaii at Manoa, Honolulu, Hawaii 96822, USA

³Department of Geology and Geophysics, Center for Computation and Technology, Louisiana State University, Baton Rouge, Louisiana 70803, USA

⁴HPCAT, Geophysical Laboratory, Carnegie Institute of Washington, Argonne, Illinois 60439, USA


⁵School of Earth and Environmental Sciences, University of Michigan, Ann Arbor, Michigan 48109, USA

*Corresponding author: Bin Chen (binchen@hawaii.edu) and Xiaojing Lai (laixiaoj@hawaii.edu)

Contents of this file

Figures S1 to S4


19
20
21



22

23 **Figure S1. Cross-section of the cell assembly for liquid structure measurements in a**
24 **Paris-Edinburgh Cell, modified after (Kono et al., 2014).** The Fe-Ni-C sample is
25 encapsulated in a boron nitride (BN) sample chamber.

26



27

28 **Figure S2. Representative experimental data and total structure factor $S(q)$ of $\text{Fe}_{90}\text{Ni}_{10}$ -**
29 **5wt.% C liquid at 1.5 GPa and 1523 K. (a)** Energy dispersive X-ray diffraction spectra of
30 $\text{Fe}_{90}\text{Ni}_{10}$ -5wt.% C liquid collected at 2θ angles from 4° to 33° . **(b)** Total structure factor $S(q)$ of
31 the $\text{Fe}_{90}\text{Ni}_{10}$ -5wt.% C liquid at 1.5 GPa and 1523 K composed of structure factor fragments from
32 measurements at different 2θ angles. The red solid curve is obtained after error-weighted
33 spline-smoothing of the overlapped structure factor fragments.




Figure S3. Additional results for the total structure factor $S(q)$ and the reduced pair distribution function $G(r)$ of Fe₉₀Ni₁₀-5wt.%C liquids at high pressures and temperatures. (a) $S(q)$, and (b) $G(r)$ of Fe₉₀Ni₁₀-5wt.%C liquids at 6.3 and 6.8 GPa.




Figure S4. Structure information of Fe₉₁Ni₉-5wt.% C liquid and Fe₃C from first-principles molecular dynamics simulations. (A) Snapshot of MD simulations with the computational super cell highlighted in dashed lines. The highlighted polyhedra are 1-atom corner-shared (red), 2-atom edge-shared (blue), 3-atom face-shared (magenta), and 4-atom distorted face-shared (green) connections of Fe/Ni – Fe/Ni pair with their second nearest neighbors. Center atoms in the polyhedra are Fe/Ni atoms, with their sizes enlarged for clarity. Fe: brown, Ni: olive, C: black. (B) Carbon coordination environment of crystalline Fe₃C. (C) Carbon coordination environment of Fe₉₁Ni₉-5wt.% C liquid at 5.1 GPa, 1673 K, the center atoms are carbon. In B, C, green polyhedral are the Voronoi polyhedral of the center carbon.